

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 069 124 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

17.01.2001 Bulletin 2001/03

(51) Int. Cl.⁷: **C07D 401/04**, C07D 401/14,
C07D 405/14, A61K 31/454,
A61P 25/04

(21) Application number: **00305981.3**

(22) Date of filing: **14.07.2000**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: **16.07.1999 WO PCT/IB99/01290**

(71) Applicant: **PFIZER INC.**

New York, N.Y. 10017 (US)

(72) Inventor: **Ito, Fumitaka**


Chita-gun, Aichi-ken 470-2393 (JP)

(74) Representative:

**Simpson, Alison Elizabeth Fraser et al
Urquhart-Dykes & Lord,
30 Welbeck Street
London W1G 8ER (GB)**

(54) **2-Benzimidazolylamine compounds as ORL1-receptor agonists**

(57) A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein

R¹ and R² are independently selected from hydrogen, halo, hydroxy, (C₁-C₄)alkyl, halo (C₁-C₄)alkyl and the like; R³ and R⁴ are independently selected from hydrogen, halo(C₁-C₁₀)alkyl, optionally substituted (C₁-C₆)alkyl and the like, or R³ and R⁴, together with the nitrogen atom to which they are attached, form an optionally substituted fully saturated, partially saturated or fully unsaturated heterocyclic ring; and R⁵ is (C₄-C₁₁)cycloalkyl and the like, has ORL1-receptor agonist activity, and are useful as analgesics or the like in mammalian subjects.

EP 1 069 124 A1

Description**Technical Field**

[0001] This invention relates to novel 1-(substituted-piperidin-4-yl)-2-benzimidazolylamine compounds, and their salts, pharmaceutical compositions containing them and their medical uses. The compounds of this invention have activity as selective ORL1-receptor agonists, and as such are useful in treating or preventing disorders or medical conditions selected from pain, inflammatory diseases and the like.

Background Art

[0002] In spite of their usefulness as analgesics, usage of opioids such as morphine and heroin are strictly limited. This is because these drugs induce side effects such as euphoria, respiratory failure, depression or constipation. Further, multiple dosage of the drugs may cause addiction. Thus, there has been a long-felt need to provide analgesics with reduced side effects.

[0003] From the above point of view, considerable pharmacological and biochemical studies have been carried out to identify opioid receptors and their endogenous ligands to prepare peptide and non-peptide opioid ligands for the receptors. In the recent past, amino acid sequences of mu- (μ -), delta (δ -) and kappa (κ -) opioid receptor subtypes have been identified and reported. Subsequently, a novel receptor subtype was identified and termed ORL1-receptor, and Meunier, J.-C *et al.* reported the isolation and structure of the endogenous agonist of the receptor (*Nature*, Vol. 377, pp. 532-535, October 12, 1995). It is suggested that the agonist compounds for ORL1-receptor be effective in neurogenic inflammation (*Tips*, Vol. 18, pp. 293-300, August 1997). It is also suggested that the agonist compounds be potent analgesics having less psychological side effects and addiction (D. Julius, *Nature*, Vol. 377, p. 476, October 12, 1995).


[0004] Neurosearch's WO 97/40035 discloses a 2-substituted 1-piperidyl benzimidazolyl compound substituted with a cycloalkyl group at the nitrogen atom of the piperidine group.

[0005] Novo Nordisk's WO 99/59997 discloses use of a small organic compound as an opioid receptor ligand for the treatment of a disease selected from migraine, non insulin dependent diabetes mellitus (type II diabetes), sepsis, inflammation, incontinence, vasomotor disturbances including the peripheral vasomotor effects such as hot flushes, and alleviating symptoms of drug withdrawal such as abstinence symptoms occurring during withdrawal from abusive drugs.

[0006] Schering Corporation's WO 00/06545 discloses use of a nociceptin receptor ORL1-agonist, alone or in combination with a second agent for treating pain, anxiety, asthma, depression, alcohol abuse, cough or allergy.

Brief Disclosure of The Invention

[0007] The present invention provides a compound of the following formula:



or a salt thereof wherein

R^1 and R^2 are independently selected from hydrogen; halo; hydroxy; (C_1-C_4) alkyl; halo (C_1-C_4) alkyl; (C_1-C_4) alkoxy; (C_1-C_4) alkylsulfonyl; (C_1-C_4) alkyl-C(=O)-; carboxy; (C_1-C_4) alkyl-C(=O)O-; amino; NH_2 -C(=O)-; (C_1-C_4) alkyl-C(=O)-NH-; (C_1-C_4) alkyl-SO₂-NH-; phenyl and naphthyl;

R^3 and R^4 are independently selected from

hydrogen;

halo(C₁-C₁₀)alkyl;

(C₁-C₆)alkyl optionally substituted with one to two substituents independently selected from hydroxy, (C₁-C₄)alkoxy, (C₁-C₄)alkyl-S-, phenoxy, amino, oxo, mono[(C₁-C₄)alkyl]amino, di[(C₁-C₄)alkyl]amino, *N*[(C₇-C₉)cycloalkyl-(C₁-C₄)alkyl]-*N*-(C₁-C₄)alkyl-amino, (C₁-C₄)alkoxy-C(=O)-, aryl selected from phenyl and naphthyl wherein the aryl is optionally substituted with one to three substituents independently selected from halo, hydroxy, (C₁-C₄)alkoxy and trifluoro(C₁-C₄)alkoxy, and heterocyclyl selected from pyrrolidyl and piperidinyl wherein the heterocyclyl is optionally substituted with (C₁-C₄)alkyl;

(C₃-C₈)cycloalkyl optionally substituted with (C₁-C₄)alkyl;

(C₃-C₈)cycloalkenyl optionally substituted with (C₁-C₄)alkoxy-C(=O)-;

5- to 6-membered-heterocyclyl, 5- to 6-membered-heterocyclyl-C(=O)- and 5- to 6-membered-heterocyclyl-(C₁-C₄)alkyl, wherein each heterocyclyl contains in the ring one to three heteroatoms independently selected from oxygen, nitrogen and sulfur, is optionally fused to a phenyl ring, and is optionally substituted with one to two substituents independently selected from halo, (C₁-C₄)alkyl, benzyl and oxo;

phenyl optionally substituted with one to three substituents independently selected from halo, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy; and

5- to 6-membered heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl contains one to four ring atoms independently selected from oxygen, nitrogen and sulfur and is optionally fused to a phenyl ring; or

R³ and R⁴, together with the nitrogen atom to which they are attached, form a fully saturated, partially saturated or fully unsaturated 5- to 8-membered nitrogen containing heterocyclic ring wherein said heterocyclic ring optionally contains in the ring one or two additional heteroatoms independently selected from nitrogen, oxygen and sulfur, optionally contains in the ring a group CR⁶R⁷ wherein R⁶ and R⁷ taken together form an oxo group or a cyclic acetal, and optionally fused to a phenyl, naphthalene or (C₅-C₈)cycloalkyl ring, and optionally substituted with one to two substituents independently selected from halo; (C₁-C₄)alkyl; halo(C₁-C₄)alkyl; (C₁-C₄)alkoxy; hydroxy; carbonyl; benzhydryl; hydroxy-(C₁-C₄)alkyl; (C₁-C₄)alkoxy-(C₁-C₄)alkyl-; amino; amino(C₁-C₄)alkyl-amino-; mercapto; (C₁-C₄)alkoxy-C(=O)-; pyrrolidino-(C₁-C₄)alkyl; amino-C(=O)-; (C₁-C₄)alkyl-C(=O)-; (C₁-C₄)alkoxy-C(=O)-amino-; piperidinyl optionally substituted with one or two substituents independently selected from amino, mono[(C₁-C₄)alkyl]amino-, di[(C₁-C₄)alkyl]amino-, benzylamino and di-(benzyl)amino; phenyl optionally substituted with one to three substituents independently selected from halo and (C₁-C₄)alkoxy, phenyl-(C₁-C₄)alkyl; phenyl-(C₁-C₄)alkenyl; phenyl-(C₁-C₄)alkoxy-C(=O)-; 1,3-benzodioxolyl-(C₁-C₄)alkyl-; trifluoromethyl; nitro; pyridyl optionally substituted with one to three substituents independently selected from halo and trifluoromethyl; quinolinyl optionally substituted with one to three substituents independently selected from halo and trifluoromethyl; pyrrolidinyl-(C₁-C₄)alkyl; and pyrrolidinyl-(CH₂)_m-CR⁸R⁹-(CH₂)_n- wherein m and n are independently 1, 2, 3 or 4, and R⁸ and R⁹, taken together with the carbon atom to which they are attached, form (C₃-C₆)cycloalkyl; and

R⁵ is phenyl or (C₄-C₁₁)cycloalkyl optionally substituted with one to three substituents independently selected from the group consisting of hydrogen, halo, hydroxy, (C₁-C₄)alkyl, halo (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylsulfonyl, (C₁-C₄)alkyl-C(=O)-, carboxy, (C₁-C₄)alkyl-C(=O)O-, amino, NH₂-C(=O)-, (C₁-C₄)alkyl-C(=O)-NH-, (C₁-C₄)alkyl-SO₂-NH-, phenyl and naphthyl.

[0008] This invention also relates to a pharmaceutical composition for the treatment of a disorder or condition mediated by ORL1-receptor and its endogenous ligands in a mammal including a human, or for anesthetizing a mammal including a human, which comprises an effective amount of the compound of formula (I) as defined above, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[0009] This invention also relates to a method of treating a disorder or condition, or anesthetizing a mammal including a human, the treatment and anesthetization of which can be effected or facilitated by agonising ORL1-receptor in a mammal including a human, comprising administering to a mammal in need of such treatment an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

Detailed Description of the Invention

[0010] The term "alkyl", as used herein, means a straight or branched saturated monovalent hydrocarbon radical including, but not limited to, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl and the like.

[0011] The term "cycloalkyl", as used herein, means a saturated carbocyclic radical including, but not limited to, cyclopropyl, cyclobutyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl and the like.

[0012] The term "alkoxy", as used herein, means an O-alkyl group wherein "alkyl" is defined above.

[0013] The term "halo", as used herein, refers to F, Cl, Br or I, preferably F or Cl.

[0014] The term "heterocyclyl", as used herein, unless otherwise indicated, includes a non-aromatic and optionally

bridged heterocyclic-group having 5 to 8 atoms comprising 1 to 4 heteroatoms each selected from oxygen (O), sulfur (S) and nitrogen (N). Examples of the heterocyclic include pyrrolidino, piperidino, morpholino, piperazinyl, homopiperidinyl and homopiperazinyl.

[0015] The term "heteroaryl", as used herein, unless otherwise indicated, refers to a monocyclic aromatic hydrocarbon group having five to six ring atoms comprising one to four heteroatoms each independently selected from N, O and S. Examples of the heteroaryl include furyl, thiophenyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, isoxazolyl, oxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl, benzofuranyl, benzothiophenyl, isoindolyl and isobenzofuranyl.

[0016] The term "treating", as used herein, refers to reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment" as used herein refers to the act of treating, as "treating" is defined immediately above.

[0017] A preferred group of compounds of the present invention includes those compounds of formula (I) wherein

R¹ and R² are independently selected from hydrogen, halo, (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₂-C₄)alkynyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy;

R³ and R⁴ are independently selected from hydrogen; halo(C₁-C₁₀)alkyl; (C₁-C₆)alkyl optionally substituted with one to two substituents independently selected from hydroxy, (C₁-C₄)alkoxy, phenoxy, amino, mono[(C₁-C₄)alkyl]amino, di[(C₁-C₄)alkyl]amino, phenyl and naphthyl; (C₃-C₈)cycloalkyl; 5- to 6-membered heterocyclyl-(C₁-C₄)alkyl wherein said heterocyclyl is selected from pyrrolidino, piperidino, morpholino, piperazinyl and homopiperazinyl, and optionally substituted with one to two substituents independently selected from halo and oxo; phenyl optionally substituted with one to three substituents independently selected from halo, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy; and heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl is selected from furyl, thiophenyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, isoxazolyl, oxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl, benzofuranyl, benzothiophenyl, isoindolyl and isobenzofuranyl; and

R⁵ is unsubstituted (C₄-C₁₁)cycloalkyl.

[0018] More preferred compounds of this invention are those compounds of formula (I), wherein

R¹ and R² are independently selected from hydrogen; halo; (C₁-C₄)alkyl; (C₂-C₄)alkenyl; (C₂-C₄)alkynyl; halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy;

R³ and R⁴ are independently selected from hydrogen; halo(C₁-C₁₀)alkyl; (C₁-C₆)alkyl optionally substituted with one to two substituents independently selected from hydroxy, (C₁-C₄)alkoxy, phenoxy, amino, mono[(C₁-C₄)alkyl]amino, di[(C₁-C₄)alkyl]amino, phenyl and naphthyl; (C₃-C₈)cycloalkyl; heterocyclyl-(C₁-C₄)alkyl wherein said heterocyclyl is selected from pyrrolidino, piperidino, morpholino, piperazinyl and homopiperazinyl, and optionally substituted with one to two substituents independently selected from halo and oxo; phenyl optionally substituted with one to three substituents independently selected from halo, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy; and heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl is selected from furyl, thiophenyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, isoxazolyl, oxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl, benzofuranyl, benzothiophenyl, isoindolyl and isobenzofuranyl; and

R⁵ is unsubstituted (C₄-C₁₁)cycloalkyl.

[0019] More preferred compounds of this invention are those compounds of formula (I), wherein

R¹ and R² are both hydrogen;

R³ and R⁴ are independently selected from hydrogen, (C₁-C₆)alkyl optionally substituted with one to three substituents independently selected from hydroxy, (C₁-C₄)alkoxy, phenoxy, di[(C₁-C₄)alkyl]amino, phenyl and naphthyl; (C₃-C₈)cycloalkyl; heterocyclyl-(C₁-C₄)alkyl wherein said heterocyclyl is selected from pyrrolidino, piperidino, morpholino and piperazinyl, and optionally substituted with one to two substituents independently selected from halo and oxo; phenyl optionally substituted with one to three halo; and heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl is selected from furyl, pyridyl and indolyl; and

R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

[0020] More preferred compounds of this invention are those compounds of formula (I), wherein R¹ and R² are both hydrogen; R³ and R⁴ are independently selected from hydrogen and (C₁-C₆)alkyl; and R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

[0021] A specific preferred compound of this invention is N-cyclohexyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1*H*-benzimidazol-2-amine, or a salt thereof.

[0022] Another preferred group of compounds of the present invention includes compounds of formula (I) wherein,

R¹ and R² are independently selected from hydrogen, halo, (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₂-C₄)alkynyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy;

R³ and R⁴, together with the nitrogen atom to which they are attached, form a 5- to 6-membered heterocyclic ring selected from pyrrolidino; piperidino; morpholino; piperazinyl and homopiperazinyl, and optionally substituted with one to two substituents independently selected from (C₁-C₄)alkyl, hydroxy, hydroxy-(C₁-C₄)alkyl, amino, amino-(C₁-C₄)alkyl, and piperidinyl optionally substituted with one or two substituents independently selected from amino, benzylamino, di(benzyl)amino, (C₁-C₄)alkyl-C(=O)- and pyrrolidino-(CH₂)_m-CR⁷R⁸-(CH₂)_n, wherein m and n are independently 1, 2, 3 or 4, and R⁷ and R⁸, taken together with the carbon atom to which they are attached, form (C₃-C₆)cycloalkyl; and

R⁵ is unsubstituted (C₄-C₁₁)cycloalkyl.

[0023] More preferred compounds of this invention are those compounds of formula (I), wherein

R¹ and R² are both hydrogen;

R³ and R⁴, together with the nitrogen atom to which they are attached, form a 5- to 6-membered heterocyclic ring selected from pyrrolidino; piperidino; morpholino; and piperazinyl optionally substituted with one to two substituents independently selected from (C₁-C₄)alkyl, hydroxy, hydroxy-(C₁-C₄)alkyl, amino, amino-(C₁-C₄)alkyl, (C₁-C₄)alkyl-C(=O)- and piperidinyl optionally substituted with one or two substituents independently selected from amino, benzylamino, di(benzyl)amino and pyrrolidino-(CH₂)_m-CR⁷R⁸-(CH₂)_n, wherein m and n are independently 1, 2, 3 or 4, and R⁷ and R⁸, taken together with the carbon atom to which they are attached, form (C₃-C₆)cycloalkyl; and

R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

[0024] More preferred compounds of this invention are those compounds of formula (I), wherein R¹ and R² are both hydrogen; R³ and R⁴, together with the nitrogen atom to which they are attached, form piperazinyl substituted with (C₁-C₄)alkyl; and R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

[0025] Specific preferred compounds of this invention include 1-(1-cyclooctylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole; 1-(1-cyclohexylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole; and 1-(1-Cycloheptylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole; or a salt thereof.

[0026] Accordingly, this invention relates to a pharmaceutical composition for the treatment of a disorder or condition mediated by ORL1-receptor and its endogenous ligands in a mammal including a human, or for anesthetizing a mammal including a human, which comprises an effective amount of the compound of formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[0027] More specifically, this invention relates to a pharmaceutical composition for the treatment of a disorder or condition selected from the group consisting of inflammatory diseases, inflammation-related hyperalgesia, eating disorders, arterial blood pressure disorders, tolerance to narcotic analgesics, dependence on narcotic analgesics, anxiety, stress disorders, psychic trauma, schizophrenia, Parkinson's disease, chorea, depressant, Alzheimer's disease, dementias, epilepsy, convulsions, migraine, non insulin dependent diabetes mellitus (type II diabetes), sepsis, incontinence, vasomotor disturbances including the peripheral vasomotor including hot flushes, alleviating symptoms of drug withdrawal including abstinence symptoms occurring during withdrawal from abusive drugs, anxiety, asthma, alcohol abuse, cough and allergy; useful as analgesics, anesthetics, neuroprotective agents or analgesic enhancers; or useful for controlling water balance, hearing regulation, controlling sodium ion excretion or ameliorating brain function, comprising an amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof that is effective in treating such disorder or condition, or effective in the use in a mammal including a human, and a pharmaceutically acceptable carrier.

[0028] This invention also relates to a method of treating a disorder or condition, or anesthetizing a mammal including a human, where the treatment or anesthetization of which can be effected or facilitated by agonising ORL1-receptor in a mammal, including a human, comprising administering to a mammal in need of such treatment an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

[0029] More specifically, this invention relates to a method for treating a disorder or condition in a mammal including a human, where the disorder or condition is selected from the group consisting of inflammatory diseases, inflammation-related hyperalgesia, eating disorder (e.g., in obesity), arterial blood pressure disorders (i.e., hypertension or hypotension), tolerance to narcotic analgesics such as morphine, dependence on narcotic analgesics such as morphine, anxiety, stress disorders, psychic trauma, schizophrenia, Parkinson's disease, chorea, depressant, Alzheimer's disease, dementias, epilepsy and convulsions, or for anesthetizing a mammal including a human, or for alleviating pain (e.g., acute, chronic and neuropathic pain), producing a neuroprotective effect, enhancing analgesic, controlling water balance (e.g., in diabetes insipidus and polyuria), hearing regulation, controlling sodium ion excretion or ameliorating brain

[0031] Examples of the second agent are antihistamines such as astemizole, azatadine,azelastin, acrivastin, brompheniramine, certirizine, cjloropheniramine, clemastine, cyclizine, carebastine, cypropheptadine, carbinoxamine, descarboethoxyloratadine, doxylamine, dimethindene, ebastine, epinastin, efletirizine, fexofenadine, hydroxyzine, keto-tifen, lorotadine, levocabastine, mizolastine, equitazine, mianserin, noberastine, meclizine, norastemizole, picumast, pyrillamine, promethazine, terfenadine, tripeleennamine, temelastine, trimeplazine and triprolidine; histamine H₃-receptor antagonists such as thioperamide, impromidine, burimamide, clobenpropit, impentamine, mifetidine, S-sopromidine, R-sopromidine, SKF-91486, GR-175737, GT-2016, UCL-1199 and clozapine; leukotriene inhibitors such as montelkast[R-(E)]-1[[[1-[3-[2-(7-chloro-2-quinolyl)-ethenyl]phenyl]-3-[2-(1-hydroxy-1-methylethyl)phenyl]propyl]thio]methyl]cyclopropaneacetic acid and its sodium salt, 1-(((R)-(3-(2-(6,7-difluoro-2-quinolinyl)ethenyl)phenyl)-3-(2-(2-hydroxy-2-propyl)phenyl)thio)methyl)cyclopropaneacetic acid and its sodium salt, 1-(((1(R)-3-(2-(2,3-dichlorothieno[3,2-b]pyridin-5-yl)-(E)-ethenyl)phenyl)-3-(2-(1-hydroxy-1-methylethyl)phenyl)propyl)thio)methyl)cyclopropaneacetic acid and its sodium salt, pranlukast, N-[4-oxo-2-(1H-tetrazol-5-yl)-4H-benzopyran-8-yl]-p-(4-phenylbutoxy)benzamide), zafirlukast, (cyclopentyl-3-[2-methoxy-4-[(o-tolylsulfonyl)carbamoyl]benzyl]-1-methylindole-5-carbamate, and [2-[[2(4-tert-butyl-2-thiazolyl)-5-benzofuranyl]oxymethyl]phenyl]acetic acid; 5-lipoxygenase inhibitors such as zileuton, docebenone, pripost, ICI-D2318 and ABT761; β-adrenergic receptor agonists such as albuterol, bitolterol, isoetharine, mataproterenol, perbuterol, salmeterol, terbutaline, isoproterenol, ephedrine and epinephrine; a xanthine derivative such as theophylline; α-adrenergic receptor agonists such as arylalkylamines (e.g., phenylpropanolamine and pseudephedrine), imidazoles (e.g., naphazoline, oxymetazoline, tetrahydrozoline and xylometazoline) and cycloalkylamines (e.g., propylhexedrine); a mast cell stabilizer such as nedocromil sodium; anti-tussive agents such as codeine, dextromethorphan, benzonatate, chlophedianol and noscapine; an expectorant such as guaifenesin; NK1-, NK2- or NK3-tachykinin receptor antagonists such as CP-99,994 and SR 48968; and GABA_B agonists such as baclofen and 3-aminopropyl-phosphinic acid.

30 **[0032]** The following reaction Schemes illustrate the preparation of the compounds of the present invention. Unless otherwise indicated R¹ to R⁸ in the reaction Schemes and discussion that follow are defined as above.

[0034] In a desired reaction step of the processes described hereafter, amino protections and removal of the amino protecting groups with reactants and reagents used may be carried out according to known procedures such as those described in *Protective Groups in Organic Synthesis* edited by T. W. Greene *et al.* (John Wiley & Sons, 1991). Typical amino protecting groups include benzyl, $C_2H_5CO_2-$ and $t\text{-ButCO}_2-$ represented as t-Boc or Boc.


40

45

50

55

Scheme 1



[0036] As shown in Scheme 1, a compound of formula (I) via a compound of formula (Ia), wherein L represents a leaving group such as halo, may be obtained from benzimidazolylpiperidine compound of formula (II) via intermediate compound of formula (IV).

[0037] First, a compound formula (II) may be subjected to a reductive amination with an aldehyde compound of formula (III) to give the compound of formula (IV). Second, the compound of formula (IV) may be reacted with a suitable nucleophilic reagent to yield the compound of formula (Ia) by introducing a leaving group to the compound of formula (IV) in the presence or absence of a catalyst. Then, the compound of formula (Ia) may be subjected to a nucleophilic aromatic substitution reaction with an amino nucleophile to give the compound of formula (I).

[0038] The reductive amination of a compound of formula (II) may be carried out using a suitable hydride reagent to give the compound of formula (IV). A suitable hydride reagent is a boron-based mild reducing reagent such as $\text{NaBH}(\text{OAc})_3$ or NaBH_3CN . This reaction can be carried out in a reaction inert solvent such as THF, $\text{ClCH}_2\text{CH}_2\text{Cl}$ or CH_2Cl_2 at about room temperature for from about 1 to about 48 hours.

[0039] The compound of formula (IV) thus obtained may be refluxed with a suitable nucleophilic reagent to give the compound of formula (Ia). In case of L is Cl, a suitable chlorinating reagent is, for example, phosphoryl chloride (POCl_3), phosphorus pentachloride (PCl_5) or phosgene (COCl_2)-triphenylphosphine (Ph_3P). In case of L is Br, a suitable bromination reagent is, for example, $\text{Br}_2\text{-Ph}_3\text{P}$. The bromination may be carried out in a reaction inert solvent such as acetonitrile or DMF. The chlorination may be carried out under conditions for example reported by R. Iemura *et al.* *J. Med. Chem.* Vol. 29, pp. 1178-1183, 1986.


[0040] The nucleophilic aromatic substitution reaction of the compound of formula (Ia) to give the compound of formula (I) may be carried out according to a known enamine formation procedure in the presence or absence of a base. Suitable bases include Hünig base (i.e., diisopropylethylamine) and inorganic bases such as K_2CO_3 . This reaction may be carried out in a reaction inert solvent at from about 0° to about 200°C (preferably from 0° to 150°C) for about 1 to about 24 hours (preferably from about 2 to about 12 hours). Suitable reaction inert solvents include alcohols such as methanol, ethanol, isopropyl alcohol, tert-butyl alcohol, and N,N-dimethylformamide (DMF) and the like. If appropriate, this reaction may be carried out in a suitable reaction chamber such as an autoclave.

[0041] A substitution reaction of a compound of formula (Ia) wherein L is Cl with an imide may also be carried out according to the procedures reported by C. H. Senanayake, *et al.*, *Tetrahedron Lett.*, Vol. 38, pp. 5607-5610, 1997. In the report, Pd-catalyst is used in the presence of a base in toluene with heating.

[0042] A compound of formula (I) may be also prepared by subjecting a compound of formula (II) according to the

preparation process illustrated in Scheme 2 comprising introduction of a leaving group, enamine formation and reductive amination.


Scheme 2



[0043] In the preparation process, first the piperidine group in a compound of formula (II) may be masked with a suitable amino-protecting group represented by Z . Suitable amino-protecting groups are benzyl-type protecting groups such as benzyl group. Benzyl group may conveniently be introduced to the compounds of formula (II) in the presence of $NaBH(OAc)_3$ or K_2CO_3 in DMF and removed by hydrogenolysis over Pd/C. Subsequent steps comprising leaving group introduction, enamine formation and reductive amination may be carried out according to the procedures illustrated in Scheme 1.

[0044] An intermediate compound of formula (IV) may be also prepared according to the procedure illustrated in Scheme 3.


Scheme 3



[0045] A compound of formula (II) may be coupled with a carbonyl chloride compound of formula (IIIb) to give the amide compound of formula (If) then reduced to the compound of formula (IV). The former coupling reaction may be carried out in the presence of a base such as triethylamine in a reaction inert solvent such as dichloromethane (CH_2Cl_2). Suitable reaction temperature ranges from 0° to $40^\circ C$, preferably at about $0^\circ C$. Suitable reaction time ranges from about 30 minutes to about 48 hours, preferably from about 1 to about 24 hours. The latter reduction may be carried out in the presence of a reducing reagent such as $LiAlH_4$ in a reaction inert solvent such as THF. Suitable reaction temperature ranges from about -78° to about $20^\circ C$, preferably from about -30° to about $0^\circ C$.

[0046] Starting from an amino-protected piperidine-4-one compound of formula (V), a compound of formula (II) may be prepared via carbonylation of an intermediate diamine compound of formula (V) according to the procedures illustrated in Scheme 4.

Scheme 4



[0047] As shown in Scheme 4, compounds of formula (II) may be prepared through the process comprising:

- reductive amination of a protected piperidine-4-one compound of formula (V) wherein Z represents an amino-protecting group to give the 4-aminopiperidine compound of formula (VI);
- coupling reaction of the compound of formula (VI) with a nitrobenzene compound of formula (VII) wherein L is a leaving group such as halo to give the nitroaniline compound of formula (VIII);
- reduction of the resulting nitroaniline compound of formula (VIII) to give the diamine compound of formula (IX);
- and
- carbonylation of the compound of formula (IX) followed by removal of the amino protecting group to give the compound of formula (II)

[0048] Each reaction step is described more specifically in the following.

- The reductive amination may be conducted by an oximation of an amino-protected piperidine-4-one compound of formula (V) followed by reduction. Both of the reactions may be conducted under conditions for oximation of carbonyl compounds known to those skilled in the art. For example, the oximation may be carried out by reacting the

piperidine compound with hydroxylamine in the presence or absence of a base in a reaction inert solvent such as alcohol at about room temperature for about 0.5 to 48 hours. The resulting oxime compound may be extracted and subjected to reduction under known conditions to give the amine compound of formula (VI). The reduction may be carried out in the presence of a reducing reagent such as lithium aluminum hydride in a reaction inert solvent such as THF at about 0 °C to room temperature for from about 0.5 to 48 hours.

(b) & (C) Steps (b) and (c) may be carried out under conditions known to those skilled in the art (e.g., B. de Costa *et al.*, *J. Chem. Soc. Perkin. Trans.*, Vol. 1, pp. 1671-1680, 1992 and N. A. Meanwell *et al.*, *Bioorganic & Medicinal Chemistry Letters*, Vol. 6, No. 14, pp. 1641-1646, 1996). For example, coupling reaction (b) may be carried out in the presence of a base such as K_2CO_3 and triethylamine (NEt_3) in a reaction inert solvent such as acetonitrile under reflux for about 0.5 to 48 hours. Then, the resulting compound of formula (VIII) may be extracted and subjected to reduction to give the diamine compound of formula (IX). The reduction may be carried out in the presence of a suitable reducing reagent such as $SnCl_2$, zinc and iron in a reaction inert solvent such as ethanol at a temperature in the range from room temperature to the reflux temperature of the reaction mixture (preferably under reflux) for from about 0.5 to about 48 hours. The reduction may also be carried out under known hydrogenation conditions such as in the presence of a metal catalyst such as Raney nickel catalysts, palladium catalysts and platinum catalysts at a temperature in the range from about 0° to 100°C (preferably at about room temperature) under hydrogen atmosphere in a reaction inert solvent such as ethanol or THF for from about 0.5 hours to 2 days.

(d) The carbonylation of the compound of formula (IX) may be carried out using a suitable carbonylating agent such as carbonyldiimidazole, trichloromethyl chloroformate, triphosgene or urea, in a reaction inert solvent such as THF, benzene, toluene or chloroform, at the temperature in the range of from about 0° to about 120°C for from about 0.5 to about 24 hours. The reaction may be conducted according to the procedures described in WO 98/54168. Then, removal of the protecting group may be carried out according to procedures known to those skilled in the art to give the compound of formula (II).

[0049] In the above preparation process, benzyl-type amino protecting groups may conveniently be introduced to or removed from the compound according to the similar conditions illustrated with Scheme 2.

[0050] Alternatively, a compound of formula (IX) may be subjected to a coupling reaction with an isothiocyanate compound and a subsequent desulfurization under known conditions to give a compound of formula (I) wherein either R^3 or R^4 is hydrogen. For example, the first coupling reaction may be carried out in a reaction inert solvent such as an alcohol (e.g., ethanol) at from about room temperature to 100° C from 30 minutes to 48 hours under stirring. The desulfurization may be carried out in the presence of an alkyl halide under reflux for from about 30 minutes to 48 hours.

[0051] A compound of formula (I) wherein $(R^3)(R^4)N$ -group has an amino or an imino group (e.g., piperidiny, piperazinyl and the like) at its terminal position may be further reacted with a desired reactant under known conditions to modify the Y. For example, these amine or imine compounds may be reacted with an alkylcarbonyl halide at about room temperature in a basic solvent to give an amide compound. The amine or imine compounds may be reacted with an amino acid, or an amino acid sulfone or sulfoxide in the presence or absence of a coupling reagent known to those skilled in the art in peptide synthesis. Suitable coupling reagents include WSC and the like. The amino or imino compound may be coupled with an amino acid, an amino acid sulfone or sulfoxide, or a phthalimido alkyl sulfonyl halide under conventional amide formation conditions in the presence of a coupling reagent in a reaction inert solvent such as acetonitrile at about room temperature. These amino acids include isoleucine, alanine, methionine, proline, phenylalanine, valine, and the like. Suitable coupling reagents are those typically used in peptide synthesis including WSC, dicyclohexylcarbodiimide (DCC), N,N' -carbonyldiimidazole (CDI), $POCl_3$, $TiCl_4$, SO_2ClF , benzotriazol-1-yl diethyl phosphate, $Ti(Obu)_4$, molecular sieves, N,N,N',N' -tetramethyl(succinimido)uronium tetrafluoroborate, CBMIT, Lawesson's reagent, chlorosulfonyl isocyanate, P_2I_4 , pyridinium salts- Bu_3N , and a mixture of Bu_3P and $PhCNO$. The amine or imine compounds may be also reacted with a guanidine compound under known conditions. A suitable reaction condition comprises reaction with an amino-protected guanidine compound in a reaction inert solvent such as THF at about room temperature (see M. S. Bernatowicz, *et al.*, *Tetrahedron Lett.*, Vol. 34,

[0052] The starting materials (III), (IIIb), (V) and (IIIV) and the other reactants are known or commercially available compounds, or may be prepared according to known procedures for a person skilled in the art.

[0053] Compounds of formula (I) of this invention may be prepared by either General Procedure A or B below.

General Procedure A:

[0054] A mixture of 2-chloro-1-(1-cyclooctylmethyl-4-piperidiny)-1*H*-benzimidazole (36 mg, 0.1 mmol) and an appropriate amine compound (about 0.6 mmol) in DMSO (about 0.5 ml) is heated at about 135°C for about 5 hrs. The volatiles are removed by N_2 blow. The residue is dissolved in CH_2Cl_2 -EtOH (about 1 ml) and loaded onto a 0.2 g/2 ml

Bond Elute (silica gel) eluting with CH_2Cl_2 . The eluent is evaporated to dryness to give the product (judged by LC-MS). If necessary, the crude product is purified by LC-MS or preparative TLC (CH_2Cl_2 -EtOH).

General Procedure B:

[0055] A mixture of 2-chlorobenzimidazole (18 mg, 0.05 mmol), an appropriate amine compound (about 0.3 mmol), and diisopropylethylamine (about 0.3 mmol) in toluene (about 0.3 ml) is heated at about 135°C for about 5 hrs. Merri-field resin and piperazinomethyl resin are added to scavenge excess reagents. The filtrate is evaporated to give the product in good yield and purity (judged by LC-MS). If necessary, the crude product is purified by LC-MS or preparative TLC.

[0056] In the each reaction described above, unless indicated otherwise, the reaction pressure is not critical. Generally, the reactions will be conducted at a pressure of about one to about three atmospheres, preferably at ambient pressure (about one atmosphere).

[0057] The subject invention also includes isotopically-labelled compounds, which are identical to those recited in formula (I), but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds of the present invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assay. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of presentation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , can afford therapeutic advantage resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirement and, hence, may be preferred in some circumstances. Isotopically labelled compounds of formula (I) of this invention and prodrugs thereof can generally, be prepared by carrying out the procedure disclosed in above-disclosed Schemes and/or Examples and Preparations below, by submitting a readily available isotopically labelled reagent for a non-isotopically labelled reagent.

[0058] The compounds of Formula (I) of this invention are basic, therefore they will form acid-addition salts. All such salts are within the scope of this invention. However, it is necessary to use an acid addition salts which is pharmaceutically-acceptable for administration to a mammal. The acid-addition salts can be prepared by standard methods. For example, the salts may be prepared by contacting the basic compounds with acid in substantially equivalent proportions in water or an organic solvent such as methanol or ethanol, or a mixture thereof. The salts can be isolated by crystallization from or evaporation of the solvent. Typical salts which can be formed are the hydrochloride, nitrate, sulfate, bisulfate, phosphate, acetate, lactate, citrate, tartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, oxalate and pamoate (1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts.

[0059] In addition, when the compounds of this invention form hydrates or solvates they are also within the scope of this invention.

[0060] The compounds of Formula (I) have been found to possess selective affinity for ORL1-receptors and ORL1 receptor agonist activity. Thus, these compounds are useful as an analgesic, anti-inflammatory, diuretic, anesthetic, neuroprotective, anti-hypertensive and anti-anxiety agent, and the like, in mammalian subjects, especially humans in need of such agents. The affinity, agonist activities and analgesic activity can be demonstrated by the following tests respectively.

Selective Affinity for ORL1-receptors:

ORL1-receptor affinity:

[0061] The ORL1 receptor binding affinity of the compounds of this invention are determined by the following procedures. Human ORL1 receptor transfected HEK-293 cell membranes and wheat-germ agglutinin coated SPA beads are combined with 0.4nM [^3H]nociceptin and unlabeled test compounds in 200 μl of 50mM Hepes buffer pH7.4 containing 10mM MgCl_2 and 1mM EDTA. This mixture is incubated at room temperature (abbreviated as rt) for 30min to 60min. Non specific binding is determined by the addition of 1 μM nociceptin. Radioactivity is counted by Wallac 1450 MicroBeta.

μ-receptor affinity:

[0062] The mu (μ) opioid receptor binding affinity of the compounds of this invention are determined by the following procedures. Human-mu opioid receptor transfected CHO-K1 cell membranes and wheat-germ agglutinin coated SPA beads are combined with 1.0nM [³H]DAMGO and unlabeled test compounds in 200μl of 50mM Hepes buffer pH7.4 containing 10mM MgCl₂ and 1mM EDTA. This mixture is incubated at rt for 30min to 60min. Non specific binding is determined by the addition of 1μM DAMGO. Radioactivity was counted by Wallac 1450 MicroBeta.

κ-receptor affinity:

[0063] The kappa (κ) opioid receptor binding affinity of the compounds of this invention are determined by the following procedures. Human kappa-opioid receptor transfected CHO-K1 cell membranes and wheat-germ agglutinin coated SPA beads are combined with 0.5nM [³H]CI-977 and unlabeled test compounds in 200μl of 50mM Hepes buffer pH7.4 containing 10mM MgCl₂ and 1mM EDTA. This mixture is incubated at rt for 30min to 60min. Non specific binding is determined by the addition of 1μM CI-977. Radio activity is counted by Wallac 1450 MicroBeta.

δ-receptor affinity:

[0064] The delta (δ) opioid receptor binding affinity of the compounds of this invention are determined by the following procedures. Human delta opioid receptor transfected CHO-K1 cell membranes and wheat-germ agglutinin coated SPA beads are combined with 2.0nM [³H]DPDPE and unlabeled test compounds in 200μl of 50mM Hepes buffer pH7.4 containing 10mM MgCl₂ and 1mM EDTA. The assay is incubated at room temperature for 30min to 60min. Non specific binding are determined by the addition of 1μM of each non-labeled ligands. Radioactivity is counted by Wallac 1450 MicroBeta.

[0065] Each percent non specific binding thus obtained is graphed as a function of compound concentration. A sigmoidal curve is used to determine 50% bindings (i.e., IC₅₀ values).

[0066] In this testing, most of the compounds prepared in the working examples appearing hereafter demonstrated higher affinity for ORL1-receptors than for mu-receptors.

$$IC_{50} \text{ (ORL1-receptors) nM} / IC_{50} \text{ (mu-receptors) nM} < 1.0$$

Functional assay:

[0067] The functional activity of the compounds of this invention in each opioid receptor can be determined in 35S-GTP S binding system according to the procedures reported by L. J. Sim, R. Xiao and S. Childers *Neuroreport* Vol. 7, pp. 729-733, 1996. Each human ORL1-, mu-, kappa- and delta- receptor transfected CHO-K1 or HEK cell membranes are used. The membranes are suspended in ice-cold 20mM HEPES buffer pH 7.4, containing 100 mM NaCl, 10 mM MgCl₂ and 1mM EDTA. 0.17 mg/ml of Dithiothreitol (DTT) is added to this buffer prior to use. Membranes are incubated at 25°C for 30 minutes with the appropriate concentration of test compounds in the presence of 5 μM GDP, 0.4 nM of 35S-GTPγS and Wheat-germ agglutinin (WGA) coated SPA bead (1.5 mg) in a 0.2 ml total volume. Basal binding is assessed in the absence of agonist, and non-specific binding is determined with 10 μM GTPγS. Radio activity is counted by Wallac 1450 MicroBeta. Some compounds of this invention prepared in Examples exhibited good ORL1 agonists activity in this assay.

Analgesic Tests:**Tail flick test:**

[0068] Male ICR mice, 4 weeks old and weighing 19-25g, are used. The training sessions are performed until mice can flick their tails within 4.0 sec by using Analgesia Meter MK-330A (Muromachi Kikai, Japan). Selected mice are used in this experiment. The latency time is recorded twice at 0.5, 1.0, and 2.0h after administration of the compound. The intensity of the beam is set to 80. Cut-off time is set to 8.0 sec. A compound of this invention is subcutaneously administered 30 min before the test. The ED₅₀ value is defined as the dose of a compound tested which halves the tail flicking observed in a control group.

Acetic acid writhing test:

[0069] Male ICR mice, 4 weeks old and weighing 21-26g, are used. They are fasted the day before use. Acetic acid

is diluted with saline to the concentration of 0.7%(v/v) and injected intraperitoneally (0.2ml/10g of body weight) to mice with a 26 gauge needle. A compound of this invention is dissolved in 0.1% methyl cellulose(MC)-saline and subcutaneously administered to mice 0.5h before acetic acid injection. After the acetic acid injection, each animal is placed in a 1L beaker and recorded by a video tape recorder. Number of writhing is counted from 5 to 15 min after acetic acid injection. The ED₅₀ value, defined as the dose of the compounds tested which halves the writhing is observed in the control group. Some compounds of this invention demonstrated good analgesic activity in this test.

Formalin licking test:

[0070] Male SD rats (80-100 g) are injected subcutaneously with a test compound dissolved in 0.1% methyl cellulose(MC)-saline or vehicle. After 30 min, 50 µl of a 2 % formalin are injected into a hind paw. The number of licking the injected paw per observation period is measured from 15 to 30 min. after the injection of formalin and expressed as % inhibition compared to the respective vehicle group. This testing method is described in, for example, (1) R.L. Follenfant, et.al., Br. J. Pharmacol. 93, 85-92 (1988); (2) H. Rogers, et.al., Br. J. Pharmacol. 106, 783-789 (1992); and (3) H. Wheeler-Aceto, et al., Psychopharmacology, 104, 35-44 (1991).

[0071] The compounds of Formula (I) of this invention can be administered by conventional pharmaceutical practice via either the oral, parenteral or topical routes to mammals, for the treatment of the indicated diseases. For administration to human patient by either route, the dosage is in the range of about 0.01mg/kg to about 3000mg/kg body weight of the patient per day, preferably about 0.01mg/kg to about 1000mg/kg body weight per day administered singly or as a divided dose. However, variations will necessarily occur depending upon the weight and condition of the subject being treated, compound employed, the disease state being treated and the particular route of administration chosen.

[0072] The compounds of the present invention may be administered alone or in combination with pharmaceutically acceptable carriers by either of the above routes previously indicated, and such administration can be carried out in single or multiple doses. Generally, the compounds can be combined with various pharmaceutically acceptable carriers in the form of tablets, powders, capsules, lozenges, trochees, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, suspensions, solutions, elixirs, syrups or the like. Such pharmaceutical carriers include solvents, excipients, coating agents, bases, binders, lubricants, disintegrants, solubilizing agents, suspending agents, emulsifying agents, stabilizers, buffering agents, tonicity agents, preservatives, flavoring agents, aromatics, coloring agents and the like.

[0073] For example, the tablets can contain various excipients such as starch, lactose, glucose, microcrystalline cellulose, calcium sulfate, calcium carbonate, talc, titanium oxide and the like, coating agents such as gelatin, hydroxypropylcellulose and the like, binding agents such as gelatin, gum arabic, methylcellulose and the like, and the disintegrating agents such as starch, agar, gelatine, sodium hydrogencarbonate and the like. Additionally, lubricating agents such as magnesium stearate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatine capsules; preferred materials in this connection also include lactose as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with diluents such as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

[0074] In general, the therapeutically-effective compounds of this invention are present in such oral dosage forms at concentration levels ranging 5% to 70% by weight, preferably 10% to 50% by weight.

[0075] The compounds of the present invention in the form of a solution may be injected parenterally such as intradermally, subcutaneously, intravenously or intramuscularly. For example the solutions are sterile aqueous solutions, aqueous suspensions and an edible oil solutions. The aqueous solutions may be suitably buffered (preferably pH>8), and may contain enough salts or glucose to make the solution isotonic with blood. The aqueous solutions are suitable for intravenous injection purposes. The aqueous suspensions may contain a suitable dispersing or suspending agents such as sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin. The aqueous suspensions can be used for subcutaneous or intramuscular injections. The edible oil such as cottonseed oil, sesame oil, coconut oil or peanut oil can be employed for the edible oil solutions. The oil solutions are suitable for intra-articular, intra-muscular and subcutaneous injection. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

[0076] It is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

[0077] For treating a respiratory disease or symptom such as cough, a compound of this invention or a salt thereof may be administered to a mammal including a human in an aerosol formulation. The formulation may be prepared according to a method known to those skilled in the art. A compound of this invention in combination with a second agent disclosed above may also be administered to a mammal including a human by a formulation described in this

specification. Those formulation may be prepared according to a method known to those skilled in the art (e.g., WO 00/06545).

Examples and Preparations

[0078] The present invention is illustrated by the following examples and preparation. However, it should be understood that the invention is not limited to the specific details of these examples and preparations. Melting points were taken with a Buchi micro melting point apparatus and is not corrected. Infrared Ray absorption spectra (IR) were measured by a Shimadzu infrared spectrometer (IR-470). ^1H and ^{13}C nuclear magnetic resonance spectra (NMR) were measured in CDCl_3 by a JEOL NMR spectrometer (JNM-GX270, 270MHz) unless otherwise indicated and peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane. The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad.

[0079] Analytical data of compounds, which can be prepared according to General Procedures A and B or were prepared in Examples hereinafter disclosed, can be taken by utilizing Waters LC-MS system (LC as 2690, ZMD as MS).

Analytical condition for LC-MS: Column YMC CombiScreen basic 4.6 mm x 50 mm, Flow rate 1 mL/min.; Mobile phase 20% MeOH/ 80% 0.1% HCO_2H in H_2O programmed over 5min to 90% MeOH/10% 0.1% HCO_2H in H_2O . Hold for 5 min.; Wave length 220-400 nm. MS detector Apcl Cone 30 Volts.

Preparation 1

2-Chloro-1-(1-cyclooctylmethyl-4-piperidinyl)-1*H*-benzimidazole

[0080] To a stirred suspension mixture of 4-(2-keto-1-benzimidazoliny)l)piperidine(17.38 g, 80 mmol) and cyclooctanecarboxaldehyde (15.39 g, 110 mmol) in CH_2Cl_2 (200 ml) was added $\text{NaBH}(\text{OAc})_3$ (21 g, 99 mmol) by portions at room temperature. After overnight reaction, the reaction mixture was diluted with CH_2Cl_2 (200 ml), washed with saturated NaHCO_3 solution and brine, and dried(Na_2SO_4). After filtration, the filtrate was concentrated to give 41.97 g of pale yellow solid. This solid was suspended in Et_2O and solid was collected byfiltration to give 24.10 g of white solid. The filtrate was concentrated to give yellow solid which was suspended in Et_2O /hexane. The solid was collected by filtration to give 2.60 g of white solid. Total 26.7 g of desired product was obtained. A mixture of this solid(26.7 g, 78.18 mmol) and POCl_3 (73 ml, 788 mmol) was refluxed for 4 days. After evaporation of excess POCl_3 , the residue was dissolved in AcOEt (200 ml). Then this solution was added dropwise to a stirred mixture of NH_4OH (200 ml of 25 % NH_4OH) and ice(100 g). The organic layer was separated and dried(Na_2SO_4). After filtration, the filtrate was concentrated to give 29.05 g of dark brown viscous oil. This oil was purified by column chromatography (silica gel: 300 g, hexane/ AcOEt : 4/1 as eluent) to afford 8.05 g of title compound and 18.56 g of the mixture included desired product mainly. This mixture was purified by column chromatography(silica gel: 300 g, hexane/ AcOEt : 4/1 as eluent) to afford 12.44 g of desired product. Total 20.49 g(72.8 %) of title compound was obtained.

^1H NMR (CDCl_3) δ 7.44-7.39(1H, m), 7.19-7.14(1H, m), 7.08(1H, dt, $J=1.3$, 7.4Hz), 6.98(1H, dt, $J=1.2$, 7.4 Hz), 4.70-4.60(1H, m), 3.76-3.28(6H, m), 2.48-1.96(4H, m), 1.86-1.25(15H, m).

Example 1

1-(1-Cyclooctylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole

[0081] A mixture of 2-chloro-1-(1-cyclooctylmethyl-4-piperidinyl)-1*H*-benzimidazole (470 mg, 1.3 mmol) and N-methylpiperazine(260 mg, 2.6 mmol) was stirred at 120C for 6 h. Then this was purified by column chromatography (silica gel: 100 g, CH_2Cl_2 /MeOH:10/1) to give 84 mg(14.9 %) of title compound.

^1H NMR (CDCl_3) δ 7.66-7.50(2H, m), 7.20-7.10(2H, m), 4.24-4.07(1H, m), 3.30-3.24(4H, m), 3.05(2H, br.d $J=11.4$ Hz), 2.66-2.60(4H, m), 2.60-2.47(2H, m), 2.38(3H, s), 2.17-1.20(21H, m).

MS(EI direct) m/z : 423(M^+), 353, 338, 206(100%), 96.

This free amine(84 mg) was treated with HCl solution in MeOH(1 ml). Then this mixture was concentrated to give yellow viscous oil which was solidified by adding Et_2O . The solid appeared was collected by filtration to afford 89 mg of light brown solid.

Anal. Calcd for $\text{C}_{26}\text{H}_{41}\text{N}_5 \cdot 3\text{HCl} \cdot 2\text{H}_2\text{O}$: C, 54.88; H, 8.50; N, 12.31. Found: C, 55.10; H, 8.82; N, 12.04.

EP 1 069 124 A1

Preparation 2

2-Chloro-1-(1-cyclohexylmethyl-4-piperidinyl)-1*H*-benzimidazole

5 [0082] This was prepared according to the procedure described in preparation 1 using cyclohexanecarboxaldehyde instead of cyclooctanecarboxaldehyde. Overall yield was 19.5 %.

¹H NMR (CDCl₃) δ 7.72-7.58(2H, m), 7.32-7.21(2H, m), 4.52-4.38(1H, m), 3.10-3.02(2H, m), 2.65-2.49(2H, m), 2.26-2.04(4H, m), 1.89-1.71(7H, m), 1.56-1.43(1H, m), 1.31-1.15(3H, m), 0.97-0.84(2H, m).

10

Example 2

1-(1-Cyclohexylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole

15 [0083] This was prepared according to the procedure described in Example 1 using 2-chloro-1-(1-cyclohexylmethyl-4-piperidinyl)-1*H*-benzimidazole instead of 2-chloro-1-(1-cyclooctylmethyl-4-piperidinyl)-1*H*-benzimidazole. Yield was 97.3 %.

¹H NMR (CDCl₃) δ 7.66-7.60(1H, m), 7.55-7.49(1H, m), 7.20-7.09(2H, m), 4.21-4.08(1H, m), 3.30-3.25(4H, m), 3.10-3.01(2H, m), 2.65-2.61(4H, m), 2.56-2.49(2H, m), 2.38(3H, s), 2.21-2.16(2H, m), 2.09-2.00(4H, m), 1.86-1.66(5H, m), 1.58-1.42(1H, m), 1.29-1.14(3H, m), 0.99-0.83(2H, m).

20

MS(ESI positive) m/z: 396(M+H)⁺.

This free amine was converted to HCl salt, mp 206-210C.

IR(KBr): 3385, 2930, 2671, 1595, 1458 cm⁻¹

25

Anal. Calcd for C₂₄H₃₇N₅·3HCl·3H₂O: C, 51.57; H, 8.29; N, 12.53. Found: C, 51.47; H, 8.62; N, 12.50.

Preparation 3

2-Chloro-1-(1-cycloheptylmethyl-4-piperidinyl)-1*H*-benzimidazole

30

[0084] This was prepared according to the procedure described in preparation 1 using cyclohepanecarboxaldehyde instead of cyclooctanecarboxaldehyde. Overall yield was 77 %.

¹H NMR (CDCl₃) δ 7.70-7.60(2H, m), 7.27-7.23(2H, m), 4.54-4.37(1H, m), 3.15-2.95(2H, m), 2.67-2.47(2H, m), 2.25-2.00(4H, m), 1.95-1.35(13H, m), 1.35-1.10(2H, m).

35

Example 3

1-(1-Cycloheptylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole

40

[0085] This was prepared according to the procedure described in Example 1 using 2-chloro-1-(1-cycloheptylmethyl-4-piperidinyl)-1*H*-benzimidazole instead of 2-chloro-1-(1-cyclooctylmethyl-4-piperidinyl)-1*H*-benzimidazole. Yield was 73 %.

¹H NMR (CDCl₃) δ 7.67-7.58(1H, m), 7.55-7.48(1H, m), 7.20-7.08(2H, m), 4.23-4.06(1H, m), 3.33-3.21(4H, m), 3.12-2.97(2H, m), 2.71-2.30(9H, m, including 3H, s, at 2.38 ppm), 2.16(2H, d, J=7.3Hz), 2.11-1.98(2H, m), 1.90-1.36(13H, m), 1.26-1.07(2H, m).

45

MS(EI direct) m/z : 409(M)⁺, 327, 227, 163, 99.

50

Preparation 4

1-(1-Cyclopentylmethyl-4-piperidinyl)-1,3-dihydro-2*H*-benzimidazole-2-one

[0086] To a stirred solution of 4-(2-keto-1-benzimidazolyl)piperidine(217.3 mg, 1 mmol) and Et₃N(15.39 g, 110 mmol) in CH₂Cl₂(5 ml) was added cyclopentanecarbonyl chloride(146 ml, 1.2 mmol) dropwise at 0C. After 2 h stirring, the reaction mixture was diluted with CH₂Cl₂, washed with saturated NaHCO₃ solution and brine, and dried(Na₂SO₄). After filtration, the filtrate was concentrated. The residue was purified by preparative TLC (1mm thick plate x 2, CH₂Cl₂/MeOH:12/1) to give 291 mg (93 %) of amide compound as white solid. To a stirred suspension of LiAlH₄(53 mg,

55

EP 1 069 124 A1

1.394 mmol) in THF(5 ml) was added a solution of the above amide derivative(291 mg, 0.929 mmol) in THF(5 ml) at -78C. The reaction mixture was stirred at -78C to -20C for 4 h. The reaction mixture was quenched with Na₂SO₄-10H₂O and diluted with CH₂Cl₂. The solid appeared was removed by filtration and the filtrate was concentrated to give oil, which was purified by preparative TLC (1mm thick plate x 2, CH₂Cl₂/MeOH:12/1) to give 202 mg (73 %) as colorless amorphous solid.

¹H NMR (CDCl₃) δ 10.30 (1H, br. s), 7.34-7.26(1H, m), 7.16-7.00(3H, m), 4.45-4.30(1H, m), 3.18-3.06(2H, m), 2.60-2.40(2H, m), 2.35(2H, d, J=7.2 Hz), 2.22-2.00(3H, m), 1.87-1.45(8H, m), 1.35-1.15(2H, m).

Example 4

1-(1-Cyclopentylmethyl-4-piperidiny)-2-(4-methylpiperaziny)-1*H*-benzimidazole

[0087] The mixture of 1-(1-cyclopentylmethyl-4-piperidiny)-1, 3-dihydro-2*H*-benzimidazole-2-one(202 mg) and POCl₃(5 ml) was heated at 120C for 2h. After cooling down to room temperature, the reaction mixture was poured into NH₄OH solution, and extracted with CH₂Cl₂. The extracts combined were washed with water, and dried (Na₂SO₄). After filtration, the filtrate was concentrated to give oil, which was purified by preparative TLC (1mm thick plate x 2, CH₂Cl₂/MeOH:15/1) to give 148 mg (69 %) of 2-chloro-1-(1-cyclopentylmethyl-4-piperidiny)-1*H*-benzimidazole as colorless amorphous solid. To a solution of 2-chloro-1-(1-cyclopentylmethyl-4-piperidiny)-1*H*-benzimidazole(148 mg, 0.466 mmol) in MeOH(5 ml) was added N-methylpiperazine(1 ml) and the resulting mixture was heated at 120C in a sealed tube for 2 days. After concentration, the residue was purified by preparative TLC (1mm thick plate x 2, CH₂Cl₂/MeOH:12/1, 2 developed) to give 161 mg of colorless amorphous solid. This was purified again by column chromatography(basic silica gel: 50 g; CH₂Cl₂/MeOH:200/1 to 50/1) to afford 52 mg (29 %) of title compound as colorless amorphous solid.

¹H NMR (CDCl₃) δ 7.66-7.58(1H, m), 7.57-7.49(1H, m), 7.20-7.08(2H, m), 4.22-4.08(1H, m), 3.32-3.24(4H, m), 3.16-3.07(2H, m), 2.66-2.30(11H, m, including 3H, s, at 2.38 ppm, and 2H, d, J=7.3Hz at 2.33 ppm), 2.18-2.00(3H, m), 1.96-1.50(8H, m), 1.34-1.16(2H, m).

MS(EI direct) m/z: 381(M)⁺, 311, 296, 217, 164(100%), 96.

Preparation 5

2-Chloro-1-(1-benzyl-4-piperidiny)-1*H*-benzimidazole

[0088] This was prepared according to the procedure described in Preparation 1 using benzaldehyde instead of cyclooctanecarboxaldehyde. Total yield was 23.1 %.

¹H NMR (CDCl₃) δ 7.70-7.60(2H, m), 7.40-7.20(7H, m), 4.52-4.42(1H, m), 3.61(2H, s), 3.20-3.07(2H, m), 2.65-2.50(2H, m), 2.25-2.15(2H, m), 1.90-1.85(2H, m).

Example 5

1-(1-Benzyl-4-piperidiny)-2-(4-methylpiperaziny)-1*H*-benzimidazole

[0089] This was prepared according to the procedure described in Example 1. Total yield was 76 %.

¹H NMR (CDCl₃) δ 7.64-7.50(2H, m), 7.40-7.10(7H, m), 4.24-4.14(1H, m), 3.59(2H, s), 3.30-3.24(4H, m), 3.12-3.05(2H, m), 2.70-2.50(6H, m), 2.38(3H, s), 2.20-2.13(2H, m), 1.80-1.75(2H, m).

MS(EI direct) m/z: 389(M⁺), 306, 217, 172, 91(100%).

Preparation 6

2-(4-Methylpiperaziny)-1-(4-piperidiny)-1*H*-benzimidazole

[0090] A suspension mixture of 1-(1-benzyl-4-piperidiny)-2-(4-methylpiperaziny)-1*H*-benzimidazole (4.62 g, 11.86 mmol) and Pd(OH)₂/C (2.35 g) in MeOH(100 ml) was stirred under hydrogen atmosphere at room temperature for 23 h. The catalyst was removed by filtration and the filtrate was concentrated to give 3.10 g (87 %) of title compound as white solid.

EP 1 069 124 A1

¹H NMR (CDCl₃) δ 7.65-7.57(2H, m), 7.20-7.10(2H, m), 4.40-4.25(1H, m), 3.50-3.40(2H, m), 3.40-3.25(4H, m), 2.90-2.82(2H, m), 2.80-2.50(7H, m), 2.40(3H, s), 1.91-1.85(2H, m).

Example 6


1-(1-Cyclobutylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole

[0091] A mixture of 2-(4-methylpiperazinyl)-1-(4-piperidinyl)-1*H*-benzimidazole (300 mg, 1 mmol), K₂CO₃ (165.9 mg, 1.2 mmol), and (bromomethyl)cyclobutane (178.8 mg, 1.2 mmol) in DMF (4 ml) was stirred at room temperature for 17 h. Then K₂CO₃ (110 mg, 0.8 mmol), and (bromomethyl)cyclobutane (119 mg, 0.8 mmol) were added to the reaction mixture. After 16 h stirring, the reaction mixture was poured into water and extracted with CH₂Cl₂. The extracts combined were dried (MgSO₄), filtered, and concentrated in vacuo to give 369 mg of oil which was purified by preparative TLC (1 mm thick plate, CH₂Cl₂/MeOH: 10/1 2 developed) to afford 147.6 mg (40 %) of title compound as light brown solid.

¹H NMR (CDCl₃) δ 7.66-7.50(2H, m), 7.20-7.08(2H, m), 4.25-4.08(1H, m), 3.35-3.20(4H, m), 3.10-2.80(3H, m), 2.70-2.45(8H, m), 2.38(3H, s), 2.20-2.05(4H, m), 2.00- 1.65(6H, m).
MS(EI direct) m/z: 367(M⁺), 341, 311, 282, 256, 217, 150(100%), 96.

[0092] The chemical structures of the compounds of Formula (I), prepared in Examples 1 to 6, are summarized in Table 1.

Table 1



| Ex. # | R ¹ | R ² | (R ³)(R ⁴)N- | R ⁵ |
|-------|----------------|----------------|--------------------------------------|----------------|
| 1 | H | H | 4-methyl-piperazin-1-yl | Cyclooctyl |
| 2 | H | H | 4-methyl-piperazin-1-yl | Cyclohexyl |
| 3 | H | H | 4-methyl-piperazin-1-yl | Cycloheptyl |
| 4 | H | H | 4-methyl-piperazin-1-yl | Cyclopentyl |
| 5 | H | H | 4-methyl-piperazin-1-yl | Phenyl |
| 6 | H | H | 4-methyl-piperazin-1-yl | Cyclobutyl |

Also, compounds of formula (I), wherein both R¹ and R² are hydrogen and R⁵ is cyclooctyl, were prepared according to General Procedure A or B described in this specification using appropriate starting materials. Those compounds are summarized in the following table.

Table 2

| Example # | General Procedure | (R ³)(R ⁴)N- | M+1 (APCI) |
|-----------|----------------------|---|---------------|
| 7 | A | Piperidino | 409 |
| 8 | A | 3-hydroxy-pyrrolidin-1-yl | 412 |
| 9 | A | 3-hydroxymethyl-piperidino | 440 |
| 10 | A | Morpholino | 412 |
| 11a | A | 4-(2-aminoethyl)-piperazin-1-yl | 454 |
| 11b | A | 2-(piperazin-1-yl)ethylamino | 454 |
| 12 | A | Cyclohexylamino | 424 |
| 13 | A | 2-(N,N-dimethylamino)ethylamino | 412 |
| 14 | A | Cyclopropylamino | 410 |
| 15 | A | (2-pyridinyl)methylamino | 432 |
| 16 | A | 3-chloroanilino | 451 |
| 17 | A | (2-furyl)methylamino | 422 |
| 18 | A | (R)-(+)-1-phenylethylamino | 446 |
| 19 | A | 2-(2-indolyl)ethylamino | 484 |
| 20 | B | (R)-(+)-2-hydroxy-1-phenylethylamino | 462 |
| 21 | B | 2-hydroxy-2-phenylethylamino | 462 |
| 22 | B | 3-(2-pyrrolidinon-1-yl)propylamino | 467 |
| 23 | B | 4-phenylpiperaziny1 | 487 |
| 24 | B | 4-acetyl-piperaziny1 | 453 |
| 25 | B | 4-[4-(N,N-dibenzylamino)-piperidino]piperidino | 688 |
| 26 | B | 1-{[1-(1-pyrrolidinylmethyl)-cyclopentyl]methyl}piperaziny1 | 576 |
| 27 | B | 2-hydroxyethylamino | 386 |
| 28 | B | 2-phenylethylamino | 446 |
| 29 | B | 3-(imidazol-1-yl)propylamino | 450 |

Table 2 - continued

| Example # | General Procedure | (R ³)(R ⁴)N- | M+1 (APCI) |
|-----------|----------------------|---|---------------|
| 30 | B | 3-methoxypropylamino | 414 |
| 31 | B | 2-phenoxyethylamino | 462 |
| 32 | B | 2-(R)-(-)-hydroxypropylamino | 400 |
| 33 | | azocan-1-yl | 437.5 |
| 34 | | pyrrolidinyl | 395.5 |
| 35 | | 2-(S)-(hydroxymethyl)pyrrolidinyl | 425.5 |
| 36 | | 4-(1,3-benzodioxol-5-ylmethyl)piperazinyl | 544.5 |
| 37 | | 1,2,3,6-tetrahydropyridin-1-yl | 407.5 |
| 38 | | 4-(2-methoxyphenyl)piperazinyl | 516.6 |
| 39 | | 4-(4-fluorophenyl)piperazinyl | 504.6 |
| 40 | | 4-benzylpiperazinyl | 500.3 |
| 41 | | 4-(2-hydroxyethyl)piperazinyl | 454.5 |
| 42 | | 1,4-dioxo-8-azaspiro[4.5]decan-8-yl | 467.5 |
| 43 | | piperidino | 409.5 |
| 44 | | 3-(ethoxycarbonyl)piperidino | 481.5 |
| 45 | | 3-methylpiperidino | 423.5 |
| 46 | | 3,5-dimethylpiperidino | 437.5 |
| 47 | | 3-(hydroxymethyl)piperidino | |
| 48 | | 4-hydroxy-4-phenylpiperidino | 501.5 |
| 49 | | 4-methylpiperidino | 423.5 |
| 50 | | 4-benzylpiperizino | |
| 51 | | decahydroisoquinolin-2-yl | 463.5 |
| 52 | | 1,2,3,4-tetrahydroisoquinolin-2-yl | 457.3 |
| 53 | | homopiperidino | 423.5 |

Table 2 - continued

| Example # | General Procedure | (R ³)(R ⁴)N- | M+1 (APCI) |
|-----------|----------------------|--|---------------|
| 54 | | 2-(S)-(methoxymethyl)pyrrolidinyl | 439.3 |
| 55 | | 2-(S)- (pyrrolidinylmethyl)pyrrolidinyl | 478.5 |
| 56 | | 4-(2-fluorophenyl)piperazinyl | 504.4 |
| 57 | | 4-methylhomopiperazinyl | 438.5 |
| 58 | | <i>trans</i> -1-cinnamylpiperazinyl | 526.5 |
| 59 | | 2-(S)-(hydroxymethyl)pyrrolidinyl | 425.5 |
| 60 | | 4- <i>t</i> -butoxycarbonylpiperazinyl | 510.4 |
| 61 | | 4-benzyloxycarbonylpiperazinyl | 544.5 |
| 62 | | 3,5-dimethylmorpholino | 439.5 |
| 63 | | 2-methyl-5-ethylpiperidino | 451.5 |
| 64 | | 4-hydroxymethylpiperidino | 439.5 |
| 65 | | 4-(2-methoxyphenyl)piperidino | 515.5 |
| 66 | | N-methyl-N-2-hydroxy-2- phenylethylamino | 475.5 |
| 67 | | 3-carbamoylpiperidino | 452.5 |
| 68 | | N-methyl-N-propylamino | 397.3 |
| 69 | | N-methyl-N-3-hydroxy-3- phenylpropylamino | 489.5 |
| 70 | | 2-(S)-methoxymethylpyrrolidinyl | |
| 71 | | 3- <i>t</i> - butoxycarbonylaminopyrrolidinyl | 510.1 |
| 72 | | N-methyl-N-(1- naphthyl)methylamino | 495.5 |
| 73 | | 4-(4-trifluoromethylpyridin-2- yl)piperazinyl | 555.5 |

Table 2 - continued

| Example # | General Procedure | (R ³)(R ⁴)N- | M+1 (APCI) |
|-----------|----------------------|--|---------------|
| 74 | | 4-(3-trifluoromethylpyridin-2-yl)piperazinyl | 555.5 |
| 75 | | 4-(3-chloro-5-trifluoromethylpyridin-2-yl)piperazinyl | 589.5 |
| 76 | | 4-(2-trifluoromethylquinolin-4-yl)piperazinyl | 605.6 |
| 77 | | 1,3,3-trimethyl-6-azabicyclo[3.2.1]octan-6-yl | 477.4 |
| 78 | | N-n-hexyl-N-methylamino | 439.4 |
| 79 | | N-2-(N,N-dimethylaminoethyl)-N-methylamino | 426.3 |
| 80 | | 3-(N,N-diethylaminocarbonyl)piperidino | 508.4 |
| 81 | | 4-(4-fluorophenyl)piperazin-1-yl | 504.4 |
| 82 | | 4-(2-nitro-4-trifluorophenyl)piperazin-1-yl | 599.4 |
| 83 | | 3-pyrrolin-1-yl | |
| 84 | | 4-formylpiperazin-1-yl | 438.4 |
| 85 | | 1,3,4,6,7,8-hexahydro-2H-pyrimido[1,2-a]pyrimidin-1-yl | 463.4 |
| 86 | | 4-acetylpiperazin-1-yl | |
| 87 | | 4-ethylpiperazin-1-yl | 438.4 |
| 88 | | 4-ethoxy-3-hydroxyethyl-piperazin-1-yl | 498.4 |

Table 2 - continued

| Example # | General Procedure | (R ³)(R ⁴)N- | M+1 (APCI) |
|-----------|----------------------|---|---------------|
| 89 | | 2,3-dihydro-1H- benzo[de]isoquinolin-2-yl | 493.4 |
| 90 | | N-(2-hydroxyethyl)-N-n-butylamino | 441.3 |
| 91 | | N, N-bis(2-pyridinylmethyl)amino | 523.4 |
| 92 | | N-(1-benzyl-3-pyrrolidinyl)-N- methylamino | 514.4 |
| 93 | | 2,6-dimethylmorpholino | 439.5 |
| 94 | | 4-(2-ethoxyphenyl)piperazin-1-yl | 530.5 |
| 95 | | 4-(3,4-dichlorophenyl)piperazin-1-yl | 554.5 |
| 96 | | 4-(3,5-dimethylphenyl)piperazin-1-yl | 514.5 |
| 97 | | 4-benzhydrylpiperazin-1-yl | 576.5 |
| 98 | | 4-(2-chlorophenyl)piperazin-1-yl | 520.5 |
| 99 | | N-cyclohexyl-N-methylamino | 437.5 |
| 100 | | N-isobutyl-N-methylamino | 411.3 |
| 101 | | N-(2-hydroxyethyl)-N-n-pentylamino | 455.4 |
| 102 | | N-(1,3-dioxolan-2-ylmethyl)-N- methylamino | 441.3 |
| 103 | | N-benzyl-N-methylamino | 445.3 |
| 104 | | 2,4-dimethyl-4,5-dihydro-1H- imidazol-1-yl | 422.3 |
| 105 | | N-(2-phenylethyl)-N-methylamino | 459.3 |
| 106 | | N-(1-methylpiperidin-4- yl)methylamino | 452.3 |

Table 2 - continued


| Example # | General Procedure | (R ³)(R ⁴)N- | M+1 (APCI) |
|-----------|----------------------|---|---------------|
| 107 | | L-prolylamino | 438.3 |
| 108 | | N-benzyl-N-2-hydroxyethylamino | 475.3 |
| 109 | | N-n-Butyl-N-4-hydroxybutylamino | 469.3 |
| 110 | | N-(2-hydroxyphenyl)methyl-N-methylamino | 461.1 |
| 111 | | 4-ethoxycarbonylpiperazin-1-yl | 482.3 |
| 112 | | N-2-(4-ethoxy-3-methoxyphenyl)ethylamino | |
| 113 | | N-3-(N,N-dimethylamino)-2,2-dimethylpropylamino | 454.3 |
| 114 | | N-3-ethoxylpropylamino | 427.3 |
| 115 | | N-(1-methyl-2-pyrrolidinyl)ethylamino | 452.3 |
| 116 | | N-4-phenylbutylamino | 473.4 |
| 117 | | N-2-(4-fluorophenyl)ethylamino | 463.3 |
| 118 | | N-ethoxycarbonyl-N-phenylamino | |
| 119 | | N-(1-ethylpyrrolidin-2-yl)methylamino | 452.3 |
| 120 | | N-[(1S,2R,5S)-6,6-dimethylbicyclo[3.1.1]hept-2-yl]methylamino | 477.4 |
| 121 | | N-(1,3-dioxolan-2-yl)ethylamino | 441.3 |
| 122 | | N-(4-methylcyclohexan-1-yl)amino | 437.5 |
| 123 | | N-(2-ethoxy)ethyl-N-methylamino | 413.3 |
| 124 | | N-3-methylcyclohexylamino | 437.3 |
| 125 | | (R)-(-)-tetrahydrofurfurylamino | 425.3 |
| 126 | | N-(2,2-dimethoxy)ethylamino | 429.2 |

Table 2 - continued

| Example # | General Procedure | (R ³)(R ⁴)N- | M+1 (APCI) |
|-----------|----------------------|---|---------------|
| 127 | | exo-2-norbornanamino | 435.3 |
| 128 | | N-(3,3-dimethylbutyl)amino | 425.3 |
| 129 | | N-(2-(S)-methylbutyl)amino | 411.3 |
| 130 | | N-2-(3,4- dimethoxyphenyl)ethylamino | 519.3 |
| 131 | | N,N-bis-2-(N',N'- diethylaminoethyl)amino | 539.3 |
| 132 | | N-2-methylthioethylamino | 415.2 |
| 133 | | N-4-trifluoromethoxybenzylamino | 515.3 |
| 134 | | N-2-methoxybenzylamino | 461.3 |
| 135 | | (3S)-2-azepanon-3-ylamino | 452.3 |
| 136 | | 3,4,5-trimethoxybenzylamino | 521.2 |
| 137 | | (1S)-1-(1naphthyl)ethylamino | 495.2 |
| 138 | | (1S)-1-(1naphthyl)ethylamino | 495.2 |
| 139 | | N-3-(2- methylpiperidino)propylamino | 480.4 |
| 140 | | (3S)-N-(1-benzylpyrrolidin-3- yl)amino | 500.4 |
| 141 | | (3R)-N-(1-benzylpyrrolidin-3- yl)amino | 500.4 |
| 142 | | N-2-(3,5- dimethoxyphenyl)ethylamino | 505.3 |
| 143 | | 3-ethoxycarbonylbicyclo[3.1.1]- heptan-2-ylamino | |
| 144 | | 6-ethoxycarbonyl-3-cyclohexen-1- ylamino | 493.3 |

Claims

1. A compound of the following formula:



or a salt thereof, wherein

R¹ and R² are independently selected from hydrogen; halo; hydroxy; (C₁-C₄)alkyl; halo (C₁-C₄)alkyl; (C₁-C₄)alkoxy; (C₁-C₄)alkylsulfonyl; (C₁-C₄)alkyl-C(=O)-; carboxy; (C₁-C₄)alkyl-C(=O)O-; amino; NH₂-C(=O)-; (C₁-C₄)alkyl-C(=O)-NH-; (C₁-C₄)alkyl-SO₂-NH-; phenyl and naphthyl;

R³ and R⁴ are independently selected from

hydrogen;

halo(C₁-C₁₀)alkyl;

(C₁-C₆)alkyl optionally substituted with one to two substituents independently selected from hydroxy, (C₁-C₄)alkoxy, (C₁-C₄)alkyl-S-, phenoxy, amino, oxo, mono[(C₁-C₄)alkyl]amino, di[(C₁-C₄)alkyl]amino, N-[(C₇-C₉)cycloalkyl-(C₁-C₄)alkyl]-N-(C₁-C₄)alkyl-amino, (C₁-C₄)alkoxy-C(=O)-, aryl selected from phenyl and naphthyl wherein the aryl is optionally substituted with one to three substituents independently selected from halo, hydroxy, (C₁-C₄)alkoxy and trifluoro(C₁-C₄)alkoxy, and heterocyclyl selected from pyrrolidyl and piperidinyl wherein the heterocyclyl is optionally substituted with (C₁-C₄)alkyl;

(C₃-C₈)cycloalkyl optionally substituted with (C₁-C₄)alkyl;

(C₃-C₈)cycloalkenyl optionally substituted with (C₁-C₄)alkoxy-C(=O)-;

5- to 6-membered-heterocyclyl, 5- to 6-membered-heterocyclyl-C(=O)- and 5- to 6-membered-heterocyclyl-(C₁-C₄)alkyl, wherein each heterocyclyl contains in the ring one to three heteroatoms independently selected from oxygen, nitrogen and sulfur, is optionally fused to a phenyl ring, and is optionally substituted with one to two substituents independently selected from halo, (C₁-C₄)alkyl, benzyl and oxo;

phenyl optionally substituted with one to three substituents independently selected from halo, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy; and

5- to 6-membered heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl contains one to four ring atoms independently selected from oxygen, nitrogen and sulfur and is optionally fused to a phenyl ring; or

R³ and R⁴, together with the nitrogen atom to which they are attached, form a fully saturated, partially saturated or fully unsaturated 5- to 8-membered nitrogen containing heterocyclic ring wherein said heterocyclic ring optionally contains in the ring one or two additional heteroatoms independently selected from nitrogen, oxygen and sulfur, optionally contains in the ring a group CR⁶R⁷ wherein R⁶ and R⁷ taken together form an oxo group or a cyclic acetal, and is optionally fused to a phenyl, naphthalene or (C₅-C₈)cycloalkyl ring, and optionally substituted with one to two substituents independently selected from halo; (C₁-C₄)alkyl; halo(C₁-C₄)alkyl; (C₁-C₄)alkoxy; hydroxy; carbonyl; benzhydryl; hydroxy-(C₁-C₄)alkyl; (C₁-C₄)alkoxy-(C₁-C₄)alkyl-; amino; amino(C₁-C₄)alkyl-amino-; mercapto; (C₁-C₄)alkoxy-C(=O)-; pyrrolidino-(C₁-C₄)alkyl; amino-C(=O)-; (C₁-C₄)alkyl-C(=O)-; (C₁-C₄)alkoxy-C(=O)-amino-; piperidinyl optionally substituted with one or two substituents independently selected from amino, mono[(C₁-C₄)alkyl]amino-, di[(C₁-C₄)alkyl]amino-, benzylamino and di-(benzyl)amino; phenyl optionally substituted with one to three substituents independently selected from halo and (C₁-C₄)alkoxy; phenyl-(C₁-C₄)alkyl; phenyl-(C₁-C₄)alkenyl; phenyl-(C₁-C₄)alkoxy-C(=O)-; 1,3-benzodioxolyl-(C₁-C₄)alkyl-; trifluoromethyl, nitro; pyridyl optionally substituted with one to three substituents independently selected from halo and trifluoromethyl; quinolinyl optionally substituted with one to three substituents independently selected from halo and trifluoromethyl; pyrrolidinyl-(C₁-C₄)alkyl; and pyrrolidinyl-(CH₂)_m-CR⁸R⁹-

(CH₂)_n- wherein m and n are independently 1, 2, 3 or 4, and R⁸ and R⁹, taken together with the carbon atom to which they are attached, form (C₃-C₆)cycloalkyl; and

R⁵ is phenyl or (C₄-C₁₁)cycloalkyl optionally substituted with one to three substituents independently selected from the group consisting of hydrogen, halo, hydroxy, (C₁-C₄)alkyl, halo (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylsulfonyl, (C₁-C₄)alkyl-C(=O)-, carboxy, (C₁-C₄)alkyl-C(=O)O-, amino, NH₂-C(=O)-, (C₁-C₄)alkyl-C(=O)-NH-, (C₁-C₄)alkyl-SO₂-NH-, phenyl and naphthyl.

2. A compound according to Claim 1, wherein

R¹ and R² are independently selected from hydrogen, halo, (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₂-C₄)alkynyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy;

R³ and R⁴ are independently selected from hydrogen; halo(C₁-C₁₀)alkyl; (C₁-C₆)alkyl optionally substituted with one to two substituents independently selected from hydroxy, (C₁-C₄)alkoxy, phenoxy, amino, mono[(C₁-C₄)alkyl]amino, di[(C₁-C₄)alkyl]amino, phenyl and naphthyl; (C₃-C₈)cycloalkyl; 5- to 6-membered heterocyclyl-(C₁-C₄)alkyl wherein said heterocyclyl is selected from pyrrolidino, piperidino, morpholino, piperazinyl and homopiperazinyl and optionally substituted with one to two substituents independently selected from halo and oxo; phenyl optionally substituted with one to three substituents independently selected from halo, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy; and heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl is selected from furyl, thiophenyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, isoxazolyl, oxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl, benzofuranyl, benzothiophenyl, isoindolyl and isobenzofuranyl; or

R³ and R⁴, together with the nitrogen atom to which they are attached, form a fully saturated, partially saturated or fully unsaturated heterocyclic ring wherein said heterocyclic ring is selected from pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, homopiperidinyl, homopiperazinyl, azocanyl, tetrahydropyridinyl, octahydroquinolyl, quinolyl, tetrahydroquinolyl, isoquinolyl, tetrahydroisoquinolyl and octahydroisoquinolyl and optionally substituted with one to two substituents independently selected from halo; (C₁-C₄)alkyl; halo(C₁-C₄)alkyl; (C₁-C₄)alkoxy; hydroxy; hydroxy-(C₁-C₄)alkyl; (C₁-C₄)alkoxy-(C₁-C₄)alkyl-; amino; amino-(C₁-C₄)alkyl-; mercapto; (C₁-C₄)alkoxy-C(=O)-; pyrrolidino-(C₁-C₄)alkyl; amino-C(=O)-; (C₁-C₄)alkyl-C(=O)-; (C₁-C₄)alkoxy-C(=O)-amino-; piperidinyl optionally substituted with one or two substituents independently selected from amino, mono[(C₁-C₄)alkyl]amino, di[(C₁-C₄)alkyl]amino-, benzylamino and di-(benzyl)amino; phenyl optionally substituted with one to three substituents selected from halo and (C₁-C₄)alkoxy; phenyl-(C₁-C₄)alkyl; phenyl-(C₁-C₄)alkenyl; phenyl-(C₁-C₄)alkoxy-C(=O)-; 1,3-benzodioxolyl-(C₁-C₄)alkyl-; pyridyl optionally substituted with one to three substituents independently selected from halo and trifluoromethyl; quinolinyl optionally substituted with one to three substituents independently selected from halo and trifluoromethyl; pyrrolidinyl-(C₁-C₄)alkyl and pyrrolidinyl-(CH₂)_m-CR⁸R⁹-(CH₂)_n-, wherein m and n are independently 1, 2, 3 or 4, and R⁸ and R⁹, taken together with the carbon atom to which they are attached, form (C₃-C₆)cycloalkyl; and

R⁵ is unsubstituted (C₄-C₁₁)cycloalkyl.

3. A compound according to Claim 2, wherein

R¹ and R² are independently selected from hydrogen, halo, (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₂-C₄)alkynyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy;

R³ and R⁴ are independently selected from hydrogen; halo(C₁-C₁₀)alkyl; (C₁-C₆)alkyl optionally substituted with one to two substituents independently selected from hydroxy, (C₁-C₄)alkoxy, phenoxy, amino, mono[(C₁-C₄)alkyl]amino, di[(C₁-C₄)alkyl]amino, phenyl and naphthyl; (C₃-C₈)cycloalkyl; heterocyclyl-(C₁-C₄)alkyl wherein said heterocyclyl is selected from pyrrolidino, piperidino, morpholino, piperazinyl and homopiperazinyl and optionally substituted with one to two substituents independently selected from halo and oxo; phenyl optionally substituted with one to three substituents independently selected from halo, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy; and heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl is selected from furyl, thiophenyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, isoxazolyl, oxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl, benzofuranyl, benzothiophenyl, isoindolyl and isobenzofuranyl; and

R⁵ is unsubstituted (C₄-C₁₁)cycloalkyl.

4. A compound according to Claim 3, wherein

R¹ and R² are both hydrogen;

R³ and R⁴ are independently selected from hydrogen; (C₁-C₆)alkyl optionally substituted with one to three substituents independently selected from hydroxy, (C₁-C₄)alkoxy, phenoxy, di[(C₁-C₄)alkyl]amino, phenyl and naphthyl; (C₃-C₈)cycloalkyl; heterocyclyl-(C₁-C₄)alkyl wherein said heterocyclyl is selected from pyrrolidino, piperidino, morpholino and piperazinyl and optionally substituted with one to two substituents independently selected from halo and oxo; phenyl optionally substituted with one to three halo; and heteroaryl-(C₁-C₄)alkyl wherein said heteroaryl is selected from furyl, pyridyl and indolyl; and

R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

5. A compound according to Claim 4, wherein R¹ and R² are both hydrogen; R³ and R⁴ are independently selected from hydrogen and (C₁-C₆)alkyl; and R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

6. A compound according to Claim 5 which is N-cyclohexyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1*H*-benzimidazol-2-amine, or a salt thereof.

7. A compound according to Claim 2, wherein

R¹ and R² are independently selected from hydrogen, halo, (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₂-C₄)alkynyl, halo(C₁-C₄)alkyl and (C₁-C₄)alkoxy;

R³ and R⁴, together with the nitrogen atom to which they are attached, form a 5- to 6-membered heterocyclic ring selected from pyrrolidino, piperidino, morpholino, piperazinyl and homopiperazinyl, and optionally substituted with one to two substituents independently selected from (C₁-C₄)alkyl, hydroxy, hydroxy-(C₁-C₄)alkyl, amino, amino-(C₁-C₄)alkyl and piperidinyl optionally substituted with one or two substituents independently selected from amino, benzylamino, di-(benzyl)amino, (C₁-C₄)alkyl-C(=O)- and pyrrolidino-(CH₂)_m-CR⁷R⁸-(CH₂)_n-, wherein m and n are independently 1, 2, 3 or 4, and R⁷ and R⁸, taken together with the carbon atom to which they are attached, form (C₃-C₆)cycloalkyl; and

R⁵ is unsubstituted (C₄-C₁₁)cycloalkyl.

8. A compound according to Claim 7, wherein

R¹ and R² are both hydrogen;

R³ and R⁴, together with the nitrogen atom to which they are attached, form a 5- to 6-membered heterocyclic ring selected from pyrrolidino, piperidino, morpholino and piperazinyl, and optionally substituted with one to two substituents independently selected from (C₁-C₄)alkyl, hydroxy, hydroxy-(C₁-C₄)alkyl, amino, amino-(C₁-C₄)alkyl, (C₁-C₄)alkyl-C(=O)- and piperidinyl optionally substituted with one or two substituents independently selected from amino, benzylamino, di(benzyl)amino and pyrrolidino-(CH₂)_m-CR⁷R⁸-(CH₂)_n-, wherein m and n are independently 1, 2, 3 or 4, and R⁷ and R⁸, taken together with the carbon atom to which they are attached, form (C₃-C₆)cycloalkyl; and

R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

9. A compound of Claim 8, wherein R¹ and R² are both hydrogen; R³ and R⁴, together with the nitrogen atom to which they are attached, form piperazinyl substituted with (C₁-C₄)alkyl; and R⁵ is unsubstituted (C₅-C₈)cycloalkyl.

10. A compound of Claim 9 selected from the group consisting of 1-(1-cyclooctylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole; 1-(1-cyclohexylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole; and 1-(1-Cycloheptylmethyl-4-piperidinyl)-2-(4-methylpiperazinyl)-1*H*-benzimidazole; or a salt thereof.

11. A pharmaceutical composition for the treatment of a disorder or condition mediated by ORL1-receptor and its endogenous ligands in a mammal including a human, or for anesthetizing a mammal including a human, which comprises an effective amount of the compound of Claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

12. A pharmaceutical composition for the treatment of a disorder or condition selected from the group consisting of inflammatory diseases, inflammation-related hyperalgesia, eating disorders, arterial blood pressure disorders, tolerance to narcotic analgesics, dependence on narcotic analgesics, anxiety, stress disorders, psychic trauma, schizophrenia, Parkinson's disease, chorea, depressant, Alzheimer's disease, dementias, epilepsy, convulsions,

migraine, non insulin dependent diabetes mellitus (type II diabetes), sepsis, incontinence, vasomotor disturbances including the peripheral vasomotor including hot flushes, alleviating symptoms of drug withdrawal including abstinence symptoms occurring during withdrawal from abusive drugs, anxiety, asthma, alcohol abuse, cough and allergy; useful as analgesics, anesthetics, neuroprotective agents or analgesic enhancers; or useful for controlling water balance, hearing regulation, controlling sodium ion excretion or ameliorating brain function, comprising an amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof that is effective in treating such disorder or condition, or effective in the use in a mammal including a human, and a pharmaceutically acceptable carrier.

13. A method of treating a disorder or condition, or anesthetizing a mammal including a human, the treatment and anesthetization of which effected or facilitated by agonising ORL1-receptor in a mammal including a human, comprising administering to a mammal in need of such treatment an effective amount of a compound of Claim 1 or a pharmaceutically acceptable salt thereof.

14. A method for treating a disorder or condition in a mammal including a human, where the disorder or condition is selected from the group consisting of inflammatory diseases, inflammation-related hyperalgesia, eating disorders, arterial blood pressure disorders, tolerance to narcotic analgesics, dependence on narcotic analgesics, anxiety, stress disorders, psychic trauma, schizophrenia, Parkinson's disease, chorea, depressant, Alzheimer's disease, dementias, epilepsy, convulsions, migraine, non insulin dependent diabetes mellitus (type II diabetes), sepsis, incontinence, vasomotor disturbances including the peripheral vasomotor including hot flushes, alleviating symptoms of drug withdrawal including abstinence symptoms occurring during withdrawal from abusive drugs, anxiety, asthma, alcohol abuse, cough and allergy; useful as analgesics, anesthetics, neuroprotective agents or analgesic enhancers; or useful for controlling water balance, hearing regulation, controlling sodium ion excretion or ameliorating brain function, comprising administering to said mammal an effective amount of a compound of Claim 1 or a pharmaceutically acceptable salt thereof that is effective in treating such disorder or condition, or effective in the use and a pharmaceutically acceptable carrier.

15. A pharmaceutical composition comprising a compound according to Claim 1 or a salt thereof in combination with a second agent for treating cough, allergy or asthma symptoms, wherein the second agent is selected from the group consisting of antihistamines; 5-lipoxygenase inhibitors; leukotriene inhibitors; H_3 inhibitors; β -adrenergic receptor agonists; xanthiane derivatives; α -adrenergic receptor agonists; mast cell stabilizers; anti-tussives; expectorants; NK1, NK2 or NK3-receptor antagonists; and GABA_B agonists.

16. A method of treating cough, allergy or asthma symptoms in a mammal including a human comprising administering to said mammal a compound according to Claim 1 or a salt thereof in combination with a second agent selected from the group consisting of antihistamines; 5-lipoxygenase inhibitors; leukotriene inhibitors; H_3 inhibitors; β -adrenergic receptor agonists; xanthiane derivatives; α -adrenergic receptor agonists; mast cell stabilizers; anti-tussives; expectorants; NK1, NK2 or NK3-receptor antagonists; and GABA_B agonists.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 00 30 5981

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|---|----------------------------------|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int.Cl.7) |
| D,X | WO 97 40035 A (NEUROSEARCH A/S) 30 October 1997 (1997-10-30) the whole document, particularly claims 11 and 13 | 1,11-14 | C07D401/04 C07D401/14 C07D405/14 A61K31/454 A61P25/04 |
| X | US 4 031 226 A (SOUDIJN W ET AL) 21 June 1977 (1977-06-21) columns 11-14, formulae XXI and XXII, and example LX | 1 | |
| Y | WO 98 54168 A (BANYU PHARMACEUTICAL CO., LTD.) 3 December 1998 (1998-12-03) * the whole document * | 1-16 | |
| P,Y | -& EP 0 990 653 A (BANYU PHARMACEUTICAL CO., LTD.) 5 April 2000 (2000-04-05) * the whole document * | 1-16 | |
| P,Y | WO 00 08013 A (PFIZER PHARMACEUTICALS INC.) 17 February 2000 (2000-02-17) * the whole document * | 1-16 | |
| D,P, Y | WO 00 06545 A (SCHERING CORPORATION) 10 February 2000 (2000-02-10) * the whole document * | 1-16 | <div>TECHNICAL FIELDS SEARCHED (Int.Cl.7)</div> C07D A61K A61P |
| INCOMPLETE SEARCH <p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>Although claims 13, 14 and 16 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.</p> | | | |
| Place of search | | Date of completion of the search | Examiner |
| THE HAGUE | | 11 October 2000 | Allard, M |
| CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document | | | |

EPO FORM 1503 03.82 (P04037)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 5981

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

11-10-2000

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9740035 A | 30-10-1997 | AU 2388697 A | 12-11-1997 |
| | | EP 0898568 A | 03-03-1999 |
| US 4031226 A | 21-06-1977 | US 3997566 A | 14-12-1976 |
| | | US 4048058 A | 13-09-1977 |
| | | US 4020097 A | 26-04-1977 |
| | | US 4118577 A | 03-10-1978 |
| | | AT 601076 A | 15-03-1981 |
| | | AU 1676276 A | 16-02-1978 |
| | | BE 845110 A | 14-02-1977 |
| | | BE 845179 A | 14-02-1977 |
| | | BG 24947 A | 15-06-1978 |
| | | CH 626337 A | 13-11-1981 |
| | | DE 2635838 A | 24-02-1977 |
| | | DE 2636614 A | 24-02-1977 |
| | | DK 365076 A | 14-02-1977 |
| | | EG 12539 A | 30-06-1979 |
| | | ES 450627 A | 01-01-1978 |
| | | ES 461191 A | 16-06-1978 |
| | | FI 762309 A | 14-02-1977 |
| | | FR 2320754 A | 11-03-1977 |
| | | FR 2353290 A | 30-12-1977 |
| | | GB 1505442 A | 30-03-1978 |
| | | GB 1552199 A | 12-09-1979 |
| | | HU 175613 B | 28-09-1980 |
| | | IE 43150 B | 31-12-1980 |
| | | IL 50252 A | 30-05-1980 |
| | | IT 1113286 B | 20-01-1986 |
| | | JP 52023080 A | 21-02-1977 |
| | | JP 1303052 C | 14-02-1986 |
| | | JP 52023064 A | 21-02-1977 |
| | | JP 60029384 B | 10-07-1985 |
| | | LU 75547 A | 25-03-1977 |
| | | NL 7608823 A | 15-02-1977 |
| | | NL 7608962 A | 15-02-1977 |
| | | NO 762781 A | 15-02-1977 |
| | | NZ 181717 A | 11-01-1979 |
| | | PH 13757 A | 17-09-1980 |
| | | PT 65471 A,B | 01-09-1976 |
| | | RO 69420 A | 30-06-1981 |
| | | SE 432928 B | 30-04-1984 |
| | | SE 7608840 A | 14-02-1977 |
| | | SE 7609024 A | 14-02-1977 |
| | | SE 8104871 A | 17-08-1981 |
| | | US 4032560 A | 28-06-1977 |
| | | US 4053485 A | 11-10-1977 |

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 5981

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

11-10-2000

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| US 4031226 A | | US 4084058 A | 11-04-1978 |
| | | US 4175202 A | 20-11-1979 |
| | | SU 694073 A | 25-10-1979 |
| | | SU 818483 A | 30-03-1981 |
| | | ZA 7604866 A | 29-03-1978 |
| | | US 4099013 A | 04-07-1978 |
| WO 9854168 A | 03-12-1998 | AU 7453298 A | 30-12-1998 |
| | | CN 1258294 T | 28-06-2000 |
| | | EP 0990653 A | 05-04-2000 |
| WO 0008013 A | 17-02-2000 | AU 4385999 A | 28-02-2000 |
| WO 0006545 A | 10-02-2000 | AU 5205699 A | 21-02-2000 |